

## RESEARCH NOTE

### Evolution, antimicrobial susceptibility and assignment to international clones of methicillin-resistant *Staphylococcus aureus* isolated over a 9-year period in two Spanish hospitals

C. Potel<sup>1</sup>, M. Álvarez<sup>1</sup>, P. Álvarez<sup>2</sup>, I. Otero<sup>1</sup>  
and E. Fluiters<sup>3</sup>

<sup>1</sup>Hospital Xeral, Microbiology Laboratory, Vigo, Pontevedra, <sup>2</sup>Complejo Hospitalario de Pontevedra, Microbiology Laboratory, Pontevedra and <sup>3</sup>Casco Vello, Primary Care, Vigo, Pontevedra, Spain

#### ABSTRACT

Antimicrobial resistance profiles, restriction fragment length polymorphism of the coagulase gene and repetitive element sequence-based PCR were used to classify 210 methicillin-resistant *Staphylococcus aureus* isolates recovered between 1997 and 2005 in two hospitals in Vigo, north-west Spain. Representative isolates belonging to the epidemic clones were analysed by *spa* typing and multilocus sequence typing, and the staphylococcal chromosomal cassette (SCC)*mec* type was determined for all isolates. The New York/Japan clone (t002-ST5-II) was detected in Spain for the first time. However, the New York/Japan and the Brazilian (t037-ST239-IIIa) clones were replaced by EMRSA-16 (t018-ST36-II), which at present is the predominant clone.

**Keywords** Epidemic clones, identification, molecular analysis, MRSA, Spain, typing

**Original Submission:** 5 September 2006; **Revised Submission:** 17 January 2007; **Accepted:** 31 January 2007

*Clin Microbiol Infect* 2007; **13**: 728–730  
10.1111/j.1469-0691.2007.01728.x

The prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) is higher in southern than in

northern Europe [1]. According to a national multicentre study, the incidence of MRSA in Spain has increased progressively [2]. During the 1980s, the Iberian clone spread throughout the country [3], although subsequent studies have demonstrated the replacement of this epidemic clone [4–6]. The aim of the present study was to identify the epidemic MRSA (EMRSA) clones in two hospitals in Vigo, north-west Spain, during the 9-year period 1997–2005. In total, 210 MRSA isolates were obtained from individual patients: 82 (39%) in internal medicine wards; 71 (33.8%) in intensive care units; 23 (11%) in trauma units; and 34 (16.2%) in other wards. The most frequent sites of isolation of MRSA were wounds (37.5%), sputum (36.6%) and intravenous catheters (11%). The frequency of MRSA in Hospital I was 28% in 1998, rising to 43% in 1999, and then decreasing to 3% in 2005, while in Hospital II the frequency increased progressively from 16% in 1997 to 22.3% in 2005. An isolate was defined as an EMRSA strain on the basis of the definition of Hoefnagels-Schuermans *et al.* [7].

Susceptibility testing was performed by the standard disk-diffusion method according to CLSI guidelines [8]. Spectinomycin susceptibility was determined as described previously [9]. The isolates were confirmed as MRSA by amplification of the *mecA* gene [10]. *S. aureus* ATCC strains 25923, BAA-40, BAA-41, BAA-43 and BAA-44 were included for quality control purposes. All isolates were screened for mupirocin resistance by the disk-diffusion method [11]; MICs were determined for isolates identified as resistant, and the presence of the *mupA* gene was determined by PCR [12]. In total, 45 isolates (15 isolates of each epidemic clone) were tested by the vancomycin agar screen method to detect heterogeneously vancomycin-resistant *S. aureus* or intermediately vancomycin-resistant *S. aureus* [13].

Analysis of restriction fragment length polymorphism of the coagulase gene (PCR-RFLP *coa*) and repetitive element sequence-based PCR (rep-PCR) were performed as described previously [14,15]. Patterns differing by one or more bands were categorised as distinct clonal types. The staphylococcal chromosome cassette (SCC)*mec* structural types were determined by a multiplex PCR strategy [16]. Randomly selected MRSA isolates belonging to the epidemic clones were further studied by *spa* typing (24 isolates) and multilocus sequence typing (six isolates) [17,18].

Corresponding author and reprint requests: C. Potel, Hospital Xeral, Microbiology Laboratory, Vigo, Pontevedra, Spain  
E-mail: carmen.potel.alvarellos@sergas.es

**Table 1.** Phenotypic and genotypic characteristics of the three major epidemic clones

Genotype: no. of isolates (%)	Resistance profile: no. of isolates (%)	SCCmec type (%)	<i>spaA</i> type (no. of isolates tested)	Sequence type (ST) (no. of isolates tested)	Clone name
A: 76 (36.2)	1: 76 (36.2) <sup>a</sup>	IIIA (36.2)	t037 (8)	ST239 (2)	Brazilian
B: 25 (11.9)	2: 22 (10.5) <sup>b</sup>	II (11.9) <sup>d</sup>	t002 (8)	ST5 (2)	New York/Japan
C: 80 (38.1)	3: 80 (38.1) <sup>c</sup>	II (38.1)	t018 (8)	ST36 (2)	EMRSA-16

<sup>a</sup>(GEN, TOB, CIP, CLI, ERY, SXT, CH)<sup>R</sup>; (RIF, TET, NIT, SPT)<sup>S</sup>.

<sup>b</sup>(CIP, CLI, ERY, SPT)<sup>R</sup>; (GEN, TOB, SXT, RIF, TET, NIT, CH)<sup>S</sup>.

<sup>c</sup>(TOB, CIP, CLI, ERY, SPT)<sup>R</sup>; (GEN, SXT, TET, NIT, CH)<sup>S</sup> (variable susceptibility to RIF).

<sup>d</sup>The band corresponding to the pUB110 insertion was not observed.


GEN, gentamicin; TOB, tobramycin; CIP, ciprofloxacin; CLI, clindamycin; ERY, erythromycin; SXT, trimethoprim-sulphamethoxazole; RIF, rifampicin; TET, tetracycline; NIT, nitrofurantoin; SPT, spectinomycin; CH, chloramphenicol.

The majority (84.8%) of MRSA isolates belonged to one of three resistance profiles, based on a panel of 11 antimicrobial agents (Table 1). None of the 45 isolates screened was intermediately vancomycin-resistant *S. aureus* or heterogeneously vancomycin-resistant *S. aureus*. Nine isolates were resistant to mupirocin by the disk-diffusion method, and three (1.4%) of these showed high-level mupirocin resistance (>256 mg/L). The *mupA* gene was only detected in these three isolates, of which two were the UK EMRSA-16 strain.

PCR-RFLP *coa*, rep-PCR and epidemiological data clustered the majority (86.2%) of the isolates into three EMRSA clonal types (Table 1). The remaining isolates (13.8%) were classified as sporadic. Overall, 95% of the isolates were hospital-acquired and 5% were associated with other healthcare facilities; no community MRSA strain was isolated.

The results of SCCmec typing, *spa* typing and multilocus sequence typing are summarised in Table 1. Fig. 1 illustrates the temporal evolution of the three EMRSA clones.

Typing is one of the most important steps involved in understanding and controlling MRSA infections. Pulsed-field gel electrophoresis analysis, although technically demanding, is considered to be the reference standard typing method [14]. However, the present study used PCR-RFLP *coa* and rep-PCR because of their accessibility for laboratories with limited resources. Moreover, these methods have been shown to have a good correlation with pulsed-field gel electrophoresis analysis [7,14]. These PCR-based methods, in conjunction with the epidemiological data, grouped the isolates into three main epidemic clones. Antimicrobial resistance patterns allowed a clear distinction among the three epidemic

**Fig. 1.** Temporal evolution of the three epidemic methicillin-resistant *Staphylococcus aureus* (MRSA) clones between 1997 and 2005 at the two hospitals included in the study.

clones and sporadic isolates, as has been reported previously [7]. Rapid classification of an isolate as epidemic or sporadic could be very useful in allowing control measures to be focused on the most important strains, thereby leading to a substantial decrease in the incidence of MRSA and the financial and logistical consequences [19].

The limited number of isolates analysed in the present study by *spa* typing and multilocus sequence typing could add some bias to the results. Nevertheless, the homogeneity in the antimicrobial resistance patterns and the results of genotyping support the decision to randomly analyse representative strains of the epidemic clones. The Iberian clone, widespread in Spain before 1996, was not detected [3,5], and the Brazilian clone predominated until 2002. The latter clone could have spread to Vigo from Portugal, where it displaced the Iberian clone during the 1990s [9]. The subsequent replacement of the multiresistant Brazilian clone by the pandemic EMRSA-16 clone suggests that EMRSA-16 could become the prevalent clone in Spanish hospitals [4,6]. The EMRSA-16 clone identified in the present study was susceptible to gentamicin but resistant to tobramycin; this phenotype has become common in French hospitals [20]. The reason(s) for the emergence and rapid spread of these strains remains unclear.

To our knowledge, this is the first report of the isolation of the New York/Japan clone in Spain. Interestingly, the plasmid pUB110 was absent from the New York/Japan isolates, which were

susceptible to gentamicin and tobramycin [16]. A similar observation has been made in Asia [21].

As in other European studies [22], no heterogeneously vancomycin-resistant *S. aureus* or intermediately vancomycin-resistant *S. aureus* isolates were identified. Nevertheless, it is necessary to continue surveillance for these phenotypes, as the use of glycopeptides continues to be high. Despite the increase in mupirocin resistance among MRSA in Spain [11], resistance to mupirocin remained at a low level (1.4%) in the present study. However, two MRSA isolates with high-level mupirocin resistance were identified as belonging to the UK EMRSA-16 clone. As the capacity of this clone to acquire resistance to mupirocin is unknown, there remains the possibility of the emergence of MRSA strains highly resistant to mupirocin in the future.

## ACKNOWLEDGEMENTS

This work was supported by grants PGIDIT05PXIB92501PR from Consellería de Innovación, Industria e Comercio and PGIDIT05SAN30PR from Consellería de Innovación, Industria e Comercio and Consellería de Sanidade, Xunta de Galicia, Spain.

## REFERENCES

1. Stefani S, Varaldo PE. Epidemiology of methicillin-resistant staphylococci in Europe. *Clin Microbiol Infect* 2003; **9**: 1179–1186.
2. Cuevas O, Cercenado E, Vindel A *et al.* Evolution of the antimicrobial resistance of *Staphylococcus* spp. in Spain: five nationwide prevalence studies, 1986–2002. *Antimicrob Agents Chemother* 2004; **48**: 4240–4245.
3. Dominguez MA, de Lencastre H, Linares J *et al.* Spread and maintenance of a dominant methicillin-resistant *Staphylococcus aureus* (MRSA) clone during an outbreak of MRSA disease in a Spanish hospital. *J Clin Microbiol* 1994; **32**: 2081–2087.
4. Perez-Roth E, Lorenzo-Díaz F, Batista N *et al.* Tracking methicillin-resistant *Staphylococcus aureus* clones during a 5-year period (1998–2002) in a Spanish hospital. *J Clin Microbiol* 2004; **42**: 4649–4656.
5. Vindel A, Trincado P, Gomez E *et al.* Prevalence and evolution of methicillin-resistant *Staphylococcus aureus* in Spanish hospitals between 1996 and 2002. *J Clin Microbiol* 2006; **44**: 266–270.
6. Montesinos I, Delgado T, Riverol D *et al.* Changes in the epidemiology of methicillin-resistant *Staphylococcus aureus* associated with the emergence of EMRSA-16 at a university hospital. *J Hosp Infect* 2006; **64**: 257–263.
7. Hoefnagels-Schuermans A, Peetermans WE, Struelens MJ *et al.* Clonal analysis and identification of epidemic strains of methicillin-resistant *Staphylococcus aureus* by antibiotyping and determination of protein A gene and coagulase gene polymorphisms. *J Clin Microbiol* 1997; **35**: 2514–2520.
8. National Committee for Clinical Laboratory Standards. *Performance standards for antimicrobial disk susceptibility tests*, 8th edn., approved standard M2-A8. Wayne, PA: NCCLS, 2003.
9. de Sousa MA, Sanches IS, Ferro ML *et al.* Intercontinental spread of a multidrug-resistant methicillin-resistant *Staphylococcus aureus* clone. *J Clin Microbiol* 1998; **36**: 2590–2596.
10. Vannuffel P, Gigi J, Ezzedine H *et al.* Specific detection of methicillin-resistant *Staphylococcus* species by multiplex PCR. *J Clin Microbiol* 1995; **33**: 2864–2867.
11. Perez-Roth E, Claverie-Martin F, Batista N *et al.* Mupirocin resistance in methicillin-resistant *Staphylococcus aureus* clinical isolates in a Spanish hospital. Co-application of multiplex PCR assay and conventional microbiology methods. *Diagn Microbiol Infect Dis* 2002; **43**: 123–128.
12. Ramsey MA, Bradley SF, Kauffman CA *et al.* Identification of chromosomal location of *mupA* gene, encoding low-level mupirocin resistance in staphylococcal isolates. *Antimicrob Agents Chemother* 1996; **40**: 2820–2823.
13. Tenover FC, Lancaster MV, Hill BC *et al.* Characterization of staphylococci with reduced susceptibilities to vancomycin and other glycopeptides. *J Clin Microbiol* 1998; **36**: 1020–1027.
14. van der Zee A, Verbakel H, van Zon JC *et al.* Molecular genotyping of *Staphylococcus aureus* strains: comparison of repetitive element sequence-based PCR with various typing methods and isolation of a novel epidemicity marker. *J Clin Microbiol* 1999; **37**: 342–349.
15. Walker J, Borrow R, Edwards-Jones V *et al.* Epidemiological characterization of methicillin-resistant *Staphylococcus aureus* isolated in the North West of England by protein A (*spa*) and coagulase (*coa*) gene polymorphisms. *Epidemiol Infect* 1998; **121**: 507–514.
16. Oliveira D, Lencastre H. Multiplex PCR strategy for rapid identification of structural types and variants of the *mec* element in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2002; **46**: 2155–2161.
17. Harmsen D, Claus H, Witte W *et al.* Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. *J Clin Microbiol* 2003; **41**: 5442–5448.
18. Enright MC, Day NP, Davies CE *et al.* Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 2000; **38**: 1008–1015.
19. Vriens M, Blok H, Fluit A *et al.* Costs associated with a strict policy to eradicate methicillin-resistant *Staphylococcus aureus* in a Dutch university medical center: a 10-year survey. *Eur J Clin Microbiol Infect Dis* 2002; **21**: 782–786.
20. Lescat C, Dupeyron C, Faubert E *et al.* Pulse-field gel electrophoresis typing of methicillin-resistant *Staphylococcus aureus* strains susceptible to aminoglycosides isolated from 1993 to 2002. *J Hosp Infect* 2004; **57**: 253–257.
21. Chongtrakool P, Ito T, Ma XX *et al.* Staphylococcal cassette chromosome *mec* (SCCmec) typing of methicillin-resistant *Staphylococcus aureus* strains isolated in 11 Asian countries: a proposal for a new nomenclature for SCCmec elements. *J Clin Microbiol* 2006; **50**: 1001–1012.
22. Borg MA, Zerafa R, Morrison D *et al.* Incidence of glycopeptide hetero-intermediate *Staphylococcus aureus* strains in Maltese hospitals. *Clin Microbiol Infect* 2005; **11**: 405–407.